

Towards a Characterisation of Pfaffian graphs

Charles H. C. Little ¹

Massey University
Palmerston North, New Zealand
e-mail: c.little@massey.ac.nz

Franz Rendl

University of Klagenfurt
Klagenfurt, Austria
e-mail: franz.rendl@uni-klu.ac.at

Ilse Fischer

University of Klagenfurt
Klagenfurt, Austria
e-mail: ilse.fischer@uni-klu.ac.at

February 1, 2008

¹The first author thanks the University of Klagenfurt for its hospitality while this research was undertaken.

Abstract

A bipartite graph G is known to be Pfaffian if and only if it does not contain an even subdivision H of $K_{3,3}$ such that $G - VH$ contains a 1-factor. However a general characterisation of Pfaffian graphs in terms of forbidden subgraphs is currently not known. In this paper we describe a possible approach to the derivation of such a characterisation. We also extend the characterisation for bipartite graphs to a slightly more general class of graphs.

1 Introduction

The graphs considered in this paper are finite and have no loops or multiple edges. They are also undirected and connected unless an indication to the contrary is given. If v and w are vertices in a directed graph, then (v, w) denotes an edge joining v and w and directed from v to w . If G is any graph, then we denote its vertex set by VG and its edge set by EG . A 1-*factor* of G is a subset f of EG such that every vertex has a unique edge of f incident on it.

Let G^* be a directed graph with an even number of vertices, and let F be the set $\{f_1, f_2, \dots, f_k\}$ of 1-factors of G^* . For all i write

$$f_i = \{(u_{i1}, w_{i1}), (u_{i2}, w_{i2}), \dots, (u_{in}, w_{in})\},$$

where $u_{ij}, w_{ij} \in VG^*$ for all j . Associate with f_i a plus sign if

$$u_{i1}w_{i1}u_{i2}w_{i2} \cdots u_{in}w_{in}$$

is an even permutation of

$$u_{11}w_{11}u_{12}w_{12} \cdots u_{1n}w_{1n},$$

and a minus sign otherwise. Thus the signs of the 1-factors are independent of the order in which their edges have been written. They are dependent on the choice of f_1 , but the resulting partition of F into two complementary subsets is not. If G is an undirected graph, we say that G is a *Pfaffian* graph if there exists a directed graph G^* with vertex set VG and edge set EG such that all the 1-factors of G^* have the same sign. We say that G^* is a *Pfaffian* orientation for G .

It is a tantalising problem to characterise Pfaffian graphs in terms of forbidden subgraphs. Pfaffian bipartite graphs have been so characterised by Little [5]. As a further contribution to a general characterisation, the present paper represents an attempt to strengthen this result.

Further progress may be possible because of a theorem of Lovász and Plummer [8, Theorem 5.4.6] on ear decompositions of 1-extendible graphs. In order to describe this theorem, we need some more definitions.

A graph is 1-*extendible* if every edge has a 1-factor containing it. We identify paths and circuits with their edge sets. If X is a path or circuit, then we denote

by VX the set of vertices of X . If P is a path and $u, v \in VP$, then we denote by $P[u, v]$ the subpath of P joining u to v . An *ear* is a path of odd cardinality. A circuit is *alternating* with respect to two given 1-factors if it is included in their symmetric difference. A circuit that is alternating with respect to a 1-factor f is also said to be *f-alternating*, or *consanguineous* (with respect to f). A path P is *f-alternating* if every internal vertex of P is incident with an edge of $P \cap f$.

Now let H be a 1-extendible subgraph of a 1-extendible graph G . Let A be an alternating circuit in G which includes $EG-EH$ and meets EH . Then an *AH-arc* (or an *HA-arc*) is a subpath of $A \cap EH$ of maximal length, and an *A $\bar{H}$ -arc* (or an *$\bar{H}A$ -arc*) is a subpath of $EG-EH$, of maximal length, whose internal vertices are in $VG-VH$. If there are n such *A $\bar{H}$ -arcs*, and each is an ear, then we say that G is obtained from H by an *n-ear adjunction*. An *ear decomposition* of G is a sequence $G_0, G_1, \dots, G_t$ of 1-extendible subgraphs of G such that G_0 is isomorphic to K_2 , $G_t = G$ and, for each $i > 0$, G_i is obtained from G_{i-1} by an *n-ear adjunction* with $n = 1$ or $n = 2$. The theorem of Lovász and Plummer alluded to earlier asserts that every 1-extendible graph has an ear decomposition. It can be stated as follows.

Theorem 1 *Let f be a 1-factor in a 1-extendible graph G . Let H be a 1-extendible proper subgraph of G such that $EH \neq \emptyset$ and $f \cap EH$ is a 1-factor of H . Then G contains an f -alternating circuit A that admits just one or two *A $\bar{H}$ -arcs*.*

In the special case where G is bipartite, it had already been shown that A may be chosen to admit just one *A $\bar{H}$ -arc*. In fact this result was implicit in [5], and was used there to prove the following characterisation of Pfaffian bipartite graphs.

Theorem 2 *A bipartite graph G is non-Pfaffian if and only if G contains an even subdivision J of $K_{3,3}$ such that $G - VJ$ has a 1-factor.*

Here we need to explain the term “even subdivision”. An *edge subdivision* of a graph G is defined as a graph obtained from G by replacing an edge joining vertices v and w with a path P joining v and w such that $VP \cap VG = \{v, w\}$. The edge subdivision is *even* if $|VP|$ is even. A graph H is a *subdivision* of G if

for some positive integer k there exist graphs $G_0, G_1, \dots, G_k$ such that $G_0 = G$, $G_k = H$ and, for all $i > 0$, G_i is an edge subdivision of G_{i-1} . If $G_1, G_2, \dots, G_k$ can be chosen so that in addition G_i is an even edge subdivision of G_{i-1} for all $i > 0$, then H is said to be an *even subdivision* of G .

The idea behind the proof of Theorem 2 runs as follows. Clearly we may assume that G is 1-extendible. If G contains J , then G is easily seen to be non-Pfaffian. Suppose on the other hand that G is non-Pfaffian. We construct an ear decomposition $G_0, G_1, \dots, G_t$ of G . Since G is bipartite, we may assume that, for each $i > 0$, G_i is obtained from G_{i-1} by the adjunction of a single ear. As G_0 is Pfaffian but G is not, there exists a smallest positive integer j such that G_j is non-Pfaffian. The graph G_j is studied in detail and eventually shown to contain J .

Theorem 1 provides a possible way to generalise this argument. If we drop the assumption that G is bipartite, then G_j is obtained from G_{j-1} by the adjunction of one or two ears. Moreover G_{j-1} might or might not be bipartite. We therefore distinguish four cases:

- (a) a 1-ear adjunction to a bipartite graph;
- (b) a 2-ear adjunction to a bipartite graph;
- (c) a 1-ear adjunction to a non-bipartite graph;
- (d) a 2-ear adjunction to a non-bipartite graph.

In each case the goal is to find a member of a class of forbidden graphs as a subgraph of G_j . In case (a), where G_j is obtained from the bipartite graph G_{j-1} by the adjunction of a single ear, this goal was achieved in [5], by showing that G_j contains an even subdivision of $K_{3,3}$. The remaining cases are still open.

It turns out that it is unnecessary to investigate case (d). In order to explain the reason, we need first to provide an account of the brick decomposition procedure of Edmonds, Lovász and Pulleyblank [2]. A graph G is *bicritical* if $G - \{u, v\}$ has a 1-factor for any pair of vertices u and v . A *brick* is a 3-connected bicritical graph.

Suppose that G is a graph which is 1-extendible but not bicritical. Then (see [8, Theorem 5.2.2(d)]) it has a maximal set S of vertices such that $|S| \geq 2$

and $G - S$ has exactly $|S|$ odd components (components with an odd number of vertices). Let $|S| = k$, and let $H_1, H_2, \dots, H_k$ be the odd components of $G - S$. For each i let G_i be the graph obtained from G by contraction of the subgraph $G - VH_i$. Let G_0 be the bipartite graph obtained from G by the successive contraction of H_i for each i . We call G_0 the *frame*. It is shown in [8, Theorem 5.2.6] that $G_0, G_1, \dots, G_k$ are 1-extendible. It is also clear that, for any i , the graph obtained from G by the contraction of H_i is 1-extendible.

Now we discard G_0 and any of $G_1, G_2, \dots, G_k$ that are isomorphic to K_2 , and we file those of $G_1, G_2, \dots, G_k$ that are bicritical. The procedure is then repeated recursively for the remaining graphs. Eventually a family of bicritical graphs is obtained.

A bicritical graph may be decomposed into bricks as follows. Let G be bicritical but not a brick. It follows that the connectivity of G is 2. Hence there are vertices u, v such that $G - \{u, v\}$ is not connected. Let $G'_1, G'_2, \dots, G'_l$ be the components of $G - \{u, v\}$, and for each i let G_i be the graph obtained from $G[VG'_i \cup \{u, v\}]$ by adjoining an edge between u and v if they are not already adjacent. It is shown in [8, Lemma 5.2.8] that G_i is bicritical. This procedure is repeated recursively until a list of bricks is obtained.

The brick decomposition procedure described above motivates the study of two operations. Firstly, let v and w be vertices, of equal degree d , in graphs H and K respectively. Let $v_1, v_2, \dots, v_d$ be the neighbours of v and $w_1, w_2, \dots, w_d$ those of w . Let G be the graph obtained from $(H - \{v\}) \cup (K - \{w\})$ by adjoining an edge between v_i and w_i for each i . Then we say that G is formed by *splicing* H and K at v and w respectively. We call $EG - (EH \cup EK)$ the *splice* of G .

Secondly, let x and y be edges in graphs H and K respectively. Let L be the graph obtained from H and K by identifying x and y to form an edge e . Then graphs L and $L - \{e\}$ are said to be obtained from H and K by *gluing* H and K at x and y . The brick decomposition procedure shows that any 1-extendible graph may be constructed from bricks by gluing and splicing, where the graphs being glued and spliced each have more than one edge.

The effect of the operations of gluing and splicing on the Pfaffian property has been studied in [6], and the results are as follows. Splicing or gluing a non-Pfaffian graph to another graph yields a non-Pfaffian graph. Gluing Pfaffian

graphs yields a Pfaffian graph. In regard to the splicing of 1-extendible Pfaffian graphs, however, the situation is a little more complicated. Let us define a *cut* in a graph G to be a minimal set X of edges such that $G - X$ has more components than has G . An example is a splice of G . A cut X is called *tight* if $|f \cap X| = 1$ for every 1-factor f . It is shown in [6] that if G is obtained by splicing 1-extendible Pfaffian graphs, and the splice in G is tight, then G is Pfaffian. It is easily checked that, if the brick decomposition procedure is reversed, then all splices are tight.

From the discussion in the previous paragraph, we see that in order to characterise Pfaffian graphs it suffices to characterise Pfaffian bricks.

Now we introduce a theorem, due to Carvalho, Lucchesi and Murty [1], concerning the smallest number of 2-ear adjunctions in an ear decomposition. The *cycle space*, $\mathcal{C}(G)$, of a graph G is the vector space spanned by the circuits of G , where the sum of vectors is defined as their symmetric difference. The *alternating space*, $\mathcal{A}(G)$, of G is the subspace of $\mathcal{C}(G)$ spanned by the alternating circuits. Note that the total number of ears adjoined in the course of an ear decomposition of a 1-extendible graph G is $\dim \mathcal{C}(G)$, for if G is obtained from a subgraph H by an n -ear adjunction then

$$\dim \mathcal{C}(G) - \dim \mathcal{C}(H) = n.$$

On the other hand,

$$\dim \mathcal{A}(G) - \dim \mathcal{A}(H) \geq 1.$$

These results imply that a lower bound for the number of 2-ear adjunctions in an ear decomposition is $\dim \mathcal{C}(G) - \dim \mathcal{A}(G)$. The theorem of Carvalho, Lucchesi and Murty alluded to earlier is that this lower bound can always be met.

Theorem 3 *The minimum number of 2-ear adjunctions in an ear decomposition of a 1-extendible graph G is $\dim \mathcal{C}(G) - \dim \mathcal{A}(G)$.*

In the case of a bipartite graph, it has been shown that no 2-ear adjunctions are necessary. It follows that if G is 1-extendible and bipartite, then $\mathcal{A}(G) = \mathcal{C}(G)$. In fact, if $G_0, G_1, \dots, G_t$ is an ear decomposition of a graph G , then G is bipartite if and only if, for each $i > 0$, G_i is obtained from G_{i-1} by the adjunction of a single ear.

The value of $\dim \mathcal{C}(G) - \dim \mathcal{A}(G)$ has been given by Lovász [7]. In the case where G is a brick, this number is 2 if G is isomorphic to the Petersen graph and 1 if G is any other brick. After noting that the Petersen graph is non-Pfaffian, we can therefore focus our attention on 1-extendible graphs G for which

$$\dim \mathcal{C}(G) - \dim \mathcal{A}(G) = 1.$$

By Theorem 3, there is an ear decomposition $G_0, G_1, \dots, G_t$ of G for which there is a unique $j > 0$ such that G_j is obtained from G_{j-1} by a 2-ear adjunction. Then G_{j-1} is bipartite but G_j is not, since j is the smallest positive integer such that G_j is obtained from G_{j-1} by a 2-ear adjunction. Hence G_j satisfies case (b). If $0 < i < j$, then G_i is bipartite and therefore satisfies case (a). If $j < i \leq t$, then G_i is non-bipartite and hence satisfies case (c). Thus case (d) does not need to be considered.

The present paper addresses case (b), which arises as follows. Suppose that G is a non-Pfaffian 1-extendible graph with ear decomposition $G_0, G_1, \dots, G_t$, and that t is the smallest integer j such that G_j is non-Pfaffian, and also the smallest integer j such that G_j is non-bipartite. Kasteleyn [3] has shown that the 1-factors of a directed graph all have equal sign if and only if all the alternating circuits are clockwise odd. (The clockwise parity of a circuit of even length is the parity of the number of its edges that are directed in agreement with a specified sense.) Since G_{t-1} is Pfaffian, we may assume it to be oriented so that each of its alternating circuits is clockwise odd. Now extend this orientation to G_t . Since G_t is non-Pfaffian, there must be alternating circuits A and B in G_t of opposite clockwise parity. As G_{t-1} is bipartite but G_t is not, G_t is obtained from G_{t-1} by a 2-ear adjunction. We show that A and B may be chosen so that each traverses both ears and there are just one or two $\bar{A}B$ -arcs. We then proceed to consider the former case. A sufficient condition for this case to arise is that $G[C \cup D]$ be bipartite whenever C and D are alternating circuits that traverse both ears and have the property that there are just two $\bar{C}D$ -arcs. Hence in this paper we confine our attention to graphs that satisfy this condition.

It is shown in [5] that $K_{3,3}$ is non-Pfaffian. It follows that no even subdivision of $K_{3,3}$ is Pfaffian. Our first lemma shows that a graph G is non-Pfaffian if it has a circuit X , of odd length, such that the graph obtained from G by contracting VX

is an even subdivision of $K_{3,3}$. In general, let us say that a graph G is *reducible* to a graph H if G has a circuit X , of odd length, such that H is obtained from G by contracting VX . Thus any graph that is reducible to an even subdivision of $K_{3,3}$ is non-Pfaffian. In fact, a graph G must be non-Pfaffian if it has a subgraph H that is reducible to an even subdivision of $K_{3,3}$ and has the property that $G - VH$ has a 1-factor.

Now let G be a 1-extendible graph that can be obtained from a 1-extendible bipartite graph by a 2-ear adjunction and suppose that $G[C \cup D]$ is bipartite whenever C and D are alternating circuits that traverse both ears and have the property that there are just two $\bar{C}D$ -arcs. In this paper we shall show that G is non-Pfaffian if and only if it has a subgraph H , reducible to an even subdivision of $K_{3,3}$, such that $G - VH$ has a 1-factor.

2 Preliminary Lemmas

A set S of alternating circuits in a directed graph G is called *intractable* if each edge of G belongs to an even number of alternating circuits in S and an odd number of the members of S are clockwise even. The former property implies that the latter is independent of the orientation of G . It is shown in [4] that G is Pfaffian if and only if it has no intractable set of alternating circuits. In fact it suffices to consider only sets of alternating circuits that are consanguineous with respect to a fixed 1-factor.

Lemma 1 *Let G be a graph with a circuit X of odd length. Let H be the graph obtained from G by contracting VX . If H is not Pfaffian, then neither is G .*

Proof. Let v be the vertex VX in H , and let $e_0, e_1, \dots, e_l$ be the edges incident on it. Since H is non-Pfaffian, it has a 1-factor f such that there is an intractable set S of f -alternating circuits. We must construct an intractable set in G .

Without loss of generality, let $e_0 \in f$. Let A be an f -alternating circuit in S , and suppose that A contains edges e_0 and e_i for some $i > 0$. In G the edges e_0 and e_i are incident on two vertices, x and y respectively, of X . Let a be the edge

of X that is incident on y but not in g , where g is the unique 1-factor of G such that $g - f \subset X$. Let $X' = X - \{a\}$, and let $A' = A \cup X'[x, y]$. (In other words, A' is the path in A of even length joining x and y .) On the other hand, if $e_0 \notin A$ then define $A' = A$.

Let $S = \{S_1, S_2, \dots, S_n\}$. Then the required intractable set of g -alternating circuits in G is $\{S'_1, S'_2, \dots, S'_n\}$ where, for each i , S'_i is constructed from S_i as described above. $\square$

We shall also need the following two lemmas, which give properties concerning the structure of the sum and union, respectively, of two consanguineous alternating circuits.

Lemma 2 *Let A_1, A_2 be f -alternating circuits in a directed graph G with 1-factor f . Then A_1 and A_2 are of opposite clockwise parity if and only if $A_1 + A_2$ includes an odd number of clockwise even alternating circuits.*

Proof. Let $A_1 = f + f_1$ and $A_2 = f + f_2$ for some 1-factors f_1 and f_2 . Suppose that A_1 and A_2 are of opposite clockwise parity. Without loss of generality, let A_1 be clockwise odd and A_2 be clockwise even. Then f has the same sign as f_1 but the opposite sign from f_2 . Hence f_1 and f_2 have opposite sign. Since $A_1 + A_2 = f_1 + f_2$, it follows that $A_1 + A_2$ includes an odd number of clockwise even alternating circuits. Similarly if A_1 and A_2 have the same clockwise parity, then f_1 and f_2 have equal sign and $A_1 + A_2$ includes an even number of clockwise even alternating circuits. $\square$

Lemma 3 *Let f be a 1-factor in a 1-extendible directed graph G . Let A and B be f -alternating circuits in G , of opposite clockwise parity, containing distinct independent edges e_1 and e_2 such that $e_1 \notin f$ and $e_2 \notin f$. Suppose that $G - \{e_1\}$ and $G - \{e_2\}$ are not bipartite but that $G - \{e_1, e_2\}$ is. Then $A \cup B$ includes alternating circuits X and Y , of opposite clockwise parity and consanguineous with respect to some 1-factor that contains neither e_1 nor e_2 , such that there are just one or two XY -arcs, each XY -arc contains e_1 or e_2 and their union contains both.*

Proof. Let $G_0 = G[A \cup B]$. Since A and B are f -alternating, they are f_0 -alternating in G_0 , where f_0 is the 1-factor $f \cap (A \cup B)$ in $G_0 - \{e_1, e_2\}$. Thus G_0 is 1-extendible. Let $A_0 = A$ and $B_0 = B$.

For some $i \geq 0$, let $G_i = G[A_i \cup B_i]$ where, for some 1-factor f_i of $G_i - \{e_1, e_2\}$, the circuits A_i and B_i are f_i -alternating circuits in G_i , of opposite clockwise parity, containing e_1 and e_2 . Thus G_i is 1-extendible. By the 2-ear theorem applied to the subgraph $G[A_i]$, the graph G_i contains an f_i -alternating circuit B'_i such that there are just one or two $\bar{A}_i B'_i$ -arcs, and in the latter case there is no f_i -alternating circuit B_i^* such that there is only one $\bar{A}_i B_i^*$ -arc.

Case 1: Suppose first that there is a unique $\bar{A}_i B'_i$ -arc P_i . Let P_i join vertices u_i and v_i . Since A_i and B'_i are f_i -alternating, $A_i + B'_i$ is an alternating circuit C_i and hence of even length. The hypotheses on G show that C_i contains both e_1 and e_2 or neither, as any circuit containing only one of e_1 and e_2 must be of odd length.

Subcase 1.1: Suppose that C_i contains e_1 and e_2 . Then $e_1 \notin B'_i$ and $e_2 \notin B'_i$. Let $A_{i+1} = C_i$ and $B_{i+1} = B_i + B'_i$, so that A_{i+1} and B_{i+1} contain e_1 and e_2 . Moreover $A_{i+1} + B_{i+1} = C_i + B_i + B'_i = A_i + B_i$, and so A_{i+1} and B_{i+1} are of opposite clockwise parity. Furthermore, since $A_{i+1} = A_i + B'_i = f_i + g_i + B'_i$ and $B_{i+1} = B_i + B'_i = f_i + h_i + B'_i$ for some 1-factors g_i and h_i of G , we see that A_{i+1} and B_{i+1} are alternating with respect to the 1-factor $f_{i+1} = f_i + B'_i$, which contains neither e_1 nor e_2 . Note also that the edge of $A_i \cap B'_i$ incident on u_i belongs to f_i and therefore to neither A_{i+1} nor B_{i+1} . Thus if we define $G_{i+1} = G[A_{i+1} \cup B_{i+1}]$ then $|EG_{i+1}| < |EG_i|$.

Subcase 1.2: Suppose that C_i contains neither e_1 nor e_2 . If C_i is clockwise even, then A_i and B'_i are the required circuits X and Y , by Lemma 2. Suppose therefore that C_i is clockwise odd. Let $A_{i+1} = B'_i = A_i + C_i$, $B_{i+1} = B_i$ and $f_{i+1} = f_i$. Then A_{i+1} and B_{i+1} are f_{i+1} -alternating and contain e_1 and e_2 . Since $C_i = A_i + A_{i+1}$ and C_i is clockwise odd, it follows from Lemma 2 that A_i and A_{i+1} have the same clockwise parity. Hence A_{i+1} and B_{i+1} are of opposite clockwise parity. Moreover the edge of $C_i - P_i$ incident on u_i belongs to neither A_{i+1} nor B_{i+1} , and so $|EG_{i+1}| < |EG_i|$, where $G_{i+1} = G[A_{i+1} \cup B_{i+1}]$.

Case 2: Suppose there are two $\bar{A}_i B'_i$ -arcs, Q_i and R_i . Let q_1 and q_2 be the

ends of Q_i , and let r_1 and r_2 be the ends of R_i . If there exists a path S , included in A_i , joining q_1 and q_2 and passing through both r_1 and r_2 , then $S \cup Q_i$ is an f_i -alternating circuit which includes Q_i but not R_i . This contradiction shows that each path included in A_i and joining q_1 and q_2 must pass through exactly one of r_1 and r_2 . We may therefore assume without loss of generality that

$$B'_i = M[r_1, q_1] \cup Q_i \cup N[q_2, r_2] \cup R_i,$$

where M is the path included in A_i which joins q_1 to q_2 and passes through r_1 , and N is the path $A_i - M$. Since A_i and B_i are f_i -alternating, the circuits $Q_i \cup M$, $Q_i \cup N$, $R_i \cup M[r_1, q_1] \cup N[q_1, r_2]$ and $R_i \cup M[r_1, q_2] \cup N[q_2, r_2]$ are of odd length. Thus we may assume without loss of generality that either $e_1 \in M[r_1, q_1]$ and $e_2 \in N[q_2, r_2]$ or $\{e_1, e_2\} \cap B'_i = \emptyset$.

Once again, let C_i be the alternating circuit $A_i + B'_i$. If C_i contains e_1 and e_2 , then the argument proceeds as in Subcase 1.1. In the remaining case we have $\{e_1, e_2\} \subseteq B'_i$, and the argument proceeds as in Subcase 1.2.

By the finiteness of G , there exists j such that G_{j+1} is not defined. Then A_j and B'_j are the required circuits. $\square$

In the remainder of this section we let e_1 and e_2 be distinct independent edges in a graph G such that neither $G - \{e_1\}$ nor $G - \{e_2\}$ is bipartite but $G - \{e_1, e_2\}$ is bipartite and 1-extendible. Let f be a 1-factor of $G - \{e_1, e_2\}$. We suppose that there exists an f -alternating circuit A in G containing e_1 and e_2 , and we let $A' = A - \{e_1\}$. Let e_1 join vertices u_1 and v_1 , and let e_2 join vertices u_2 and v_2 , where $e_2 \in A'[u_1, v_2]$.

Lemma 4 *Let C be an f -alternating circuit in $G - \{e_1, e_2\}$, and let a be an edge of $A'[v_1, v_2] \cap C$ joining vertices u and v , where $a \in A'[v_2, u]$. Let $C' = C - \{a\}$. Let b be an edge of C' joining vertices w and x , where $b \in C'[v, x]$.*

- (a) *If $b \in A'[v_1, v_2]$, then $b \in A'[v_2, w]$.*
- (b) *If $b \in A'[u_1, u_2]$, then $b \in A'[u_2, w]$.*

Proof. Let X be an $\bar{A}C$ -arc joining vertices y and z . Since A and C are both f -alternating, there must be an edge $c \in f \cap A \cap C$ incident on y and an edge

$d \in f \cap A \cap C$ incident on z . If $e_2 \in A'[y, z]$ then the circuit $X \cup A'[y, z]$ contains e_2 but not e_1 , and is therefore of odd length by the hypotheses on G . In this case exactly one of c and d is in $A'[y, z]$. Similarly if $e_2 \notin A'[y, z]$ then either c and d are both in $A'[y, z]$ or neither of them is. The lemma follows by applying these observations sequentially to all the $\bar{A}C$ -arcs. $\square$

3 The Main Results

In order to prepare for the proof of our first theorem, we need the following rather technical definition.

Let G be a graph with 1-factor f . Let A be an f -alternating path in G joining vertices u and v . Let x_0 and y_0 be distinct vertices of VA , where $x_0 \in VA[y_0, v]$. Suppose that the edges of f incident on x_0 and y_0 belong to $A[x_0, v]$ and $A[u, y_0]$ respectively. Let $C_1, C_2, \dots, C_k$ be f -alternating circuits with the following properties.

- (a) For each i there exists a unique AC_i -arc. Let x_i and y_i be the ends of this arc, where $x_i \in VA[u, y_i]$.
- (b) We have $x_1 \in VA[u, y_0] - \{y_0\}$, $y_1 \in VA[y_0, v] - \{y_0\}$ and $y_k \in VA[x_0, v] - \{x_0\}$.
- (c) If $k > 1$, then for each $i \geq 2$ we have $x_i \in VA[y_{i-2}, y_{i-1}] - \{y_{i-2}, y_{i-1}\}$ and $y_i \in VA[y_{i-1}, v] - \{y_{i-1}\}$.

Then the sequence

$$(C_1, C_2, \dots, C_k)$$

is called a *cascade* of length k along $A[y_0, x_0]$ from y_0 to x_0 .

Theorem 4 *Let G be a 1-extendible graph with 1-factor f . Let e_1 and e_2 be distinct independent edges of $EG - f$ such that neither $G - \{e_1\}$ nor $G - \{e_2\}$ is bipartite but $G - \{e_1, e_2\}$ is bipartite, Pfaffian and 1-extendible. Suppose there exist f -alternating circuits A and B , both containing e_1 and e_2 , such that there*

is a unique AB -arc and $A + B$ is clockwise even under a Pfaffian orientation of $G - \{e_1, e_2\}$. Then G has a subgraph H , reducible to an even subdivision of $K_{3,3}$, such that $G - VH$ has a 1-factor.

Proof. Let e_1 join vertices u_1 and v_1 , and let $A' = A - \{e_1\}$. Let e_2 join vertices u_2 and v_2 , where $u_2 \in VA'[u_1, v_2]$. Let x_0 and y_0 be the ends of the unique $\bar{A}B$ -arc P . Since $\{e_1, e_2\} \subset A \cap B$, we may assume with no less generality that $x_0 \in VA'[v_1, v_2]$ and $y_0 \in VA'[v_2, x_0]$. Moreover the edge a of f incident on y_0 is in $A'[v_2, y_0]$ and the edge of f incident on x_0 is in $A'[x_0, v_1]$. We may assume A and B to have been chosen to maximise $|A'[v_2, y_0]| + |A'[x_0, v_1]|$. Let b be the edge of $A'[x_0, y_0]$ incident on y_0 .

Claim 1: There is a cascade $\mathcal{C}$ along $A'[y_0, x_0]$ from y_0 to x_0 such that distinct elements of this cascade meet, if at all, only in $A'[x_0, y_0]$.

Let $K = G - \{e_1, e_2\}$. Since K is 1-extendible, there is an f -alternating circuit C_1 in K containing a and b . Let $C'_1 = C_1 - \{a\}$.

Since $A'[v_2, y_0] \cap C_1 \neq \emptyset$ and $A'[y_0, v_1] \cap C_1 \neq \emptyset$, we may select a vertex

$$x_1 \in (VA'[v_2, y_0] - \{y_0\}) \cap VC_1$$

that minimises $|C'_1[x_1, y_0]|$ and a vertex

$$y_1 \in VA'[y_0, v_1] \cap VC_1$$

that minimises $|C'_1[x_1, y_1]|$. We may assume that C_1 has been chosen to minimise $|A'[y_1, v_1]|$, and by Lemma 4 we may assume that $A'[x_1, y_1]$ is an AC_1 -arc.

From Lemma 4 we see that $y_1 \neq x_0$. If $y_1 \in VA'[y_0, x_0]$, then we repeat the argument. Thus there must be an f -alternating circuit C_2 in K containing both the edges of A incident on y_1 . Let $C'_2 = C_2 - \{a'\}$, where a' is the edge of f incident on y_1 . We may select a vertex

$$x_2 \in (VA'[v_2, y_1] - \{y_1\}) \cap VC_2$$

to minimise $|C'_2[x_2, y_1]|$ and a vertex

$$y_2 \in VA'[y_1, v_1] \cap VC_2$$

to minimise $|C'_2[x_2, y_2]|$. In fact it follows from the choice of C_1 that $x_2 \in VA'[y_0, y_1]$. We may assume that C_2 has been chosen to minimise $|A'[y_2, v_1]|$. By Lemma 4 we may also suppose that $A'[x_2, y_2]$ is an AC_2 -arc.

Suppose that

$$C'_1[x_1, y_1] \cap C'_2[x_2, y_2] \neq \emptyset.$$

Let w be the vertex of $VC'_1[x_1, y_1] \cap VC'_2[x_2, y_2]$ that minimises $|C'_2[w, y_2]|$. Since the circuit

$$C'_1[w, y_1] \cup A'[y_1, y_2] \cup C'_2[y_2, w]$$

contains neither e_1 nor e_2 and is therefore of even length, it follows that

$$C'_1[x_1, w] \cup C'_2[w, y_2] \cup A'[y_2, x_1]$$

is an f -alternating circuit containing a and b . Since $y_2 \in VA'[y_1, v_1] - \{y_1\}$, the choice of C_1 is contradicted. We conclude that $C'_1[x_1, y_1] \cap C'_2[x_2, y_2] = \emptyset$.

From Lemma 4 we see that $y_2 \neq x_0$. If $y_2 \in VA'[y_0, x_0]$, then we repeat the argument inductively. By the finiteness of G , there exists $k > 0$ such that $y_k \in VA'[x_0, v_1]$. Then $(C_1, C_2, \dots, C_k)$ is the required cascade $\mathcal{C}$ along $A'[y_0, x_0]$ from y_0 to x_0 . The proof of the claim is now complete.

For each i , let p_i be the number of BC_i -arcs included in P , and let q_i be the number of AC_i -arcs. We may assume f , A , B and $\mathcal{C}$ chosen to minimise

$$\sum_{i=1}^k (p_i + q_i). \quad (1)$$

Since distinct elements of $\mathcal{C}$ meet, if at all, only in $A'[x_0, y_0]$, it follows that

$$\bigcup_{i=1}^{k-1} (C'_i[x_i, y_i] \cup A'[y_i, x_{i+1}]) \cup C'_k[x_k, y_k]$$

is a path X joining x_1 to y_k . Traversed from x_1 to y_k , it meets a succession of AC_i -arcs and BC_i -arcs. A *trace* of $\mathcal{C}$ is the sequence obtained from X by recording 0 for each AC_i -arc included in $A'[x_0, y_0]$, 1 for each BC_i -arc included in P and 2 for each AC_i -arc included in $A'[u_1, u_2]$. (This sequence is not uniquely determined unless A and B have been specified.) The next two claims show that, up to homeomorphism, the graph

$$G[A \cup B \cup \bigcup_{i=1}^k C_i]$$

is determined by a trace of $\mathcal{C}$.

Claim 2: Suppose that for some i and $j > i$ there are a BC_i -arc included in P and a BC_j -arc included in P . Let the former arc join vertices u and v , where $u \in VC'_i[x_i, v]$, and the latter arc vertices w and x , where $w \in VC'_j[x_j, x]$. Then $x \in VP[w, y_0]$, $u \in VP[x, y_0]$ and $v \in VP[u, y_0]$.

By Lemma 4 we see that $v \in VP[u, y_0]$ and $x \in VP[w, y_0]$. We must show that $u \in VP[x, y_0]$. Suppose therefore that $w \in VP[v, y_0]$. We may suppose u, v, w, x to have been chosen to minimise $|P[v, w]|$. It follows that the circuit

$$C'_i[x_i, v] \cup P[v, w] \cup C'_j[w, y_j] \cup A'[y_j, x_i]$$

contradicts the choice of C_i . The claim has now been verified.

The same argument can also be applied to AC_i -arcs and AC_j -arcs meeting $A'[u_1, u_2]$. In other words, suppose that for some i and $j > i$ there are an AC_i -arc meeting $A'[u_1, u_2]$ and an AC_j -arc meeting $A'[u_1, u_2]$. Let the former arc join vertices u and v , where $u \in VC'_i[x_i, v]$, and the latter arc vertices w and x , where $w \in VC'_j[x_j, x]$. Then $x \in VA'[w, u_2]$, $u \in VA'[x, u_2]$ and $v \in VA'[u, u_2]$.

Claim 3: Suppose that for some i there are two BC_i -arcs included in P . Let one of these arcs join vertices u and v , where $u \in VC'_i[x_i, v]$, and let the other join vertices w and x , where $w \in VC'_i[v, y_i]$ and $x \in VC'_i[w, y_i]$. Then $x \in VP[w, y_0]$, $u \in VP[x, y_0]$ and $v \in VP[u, y_0]$.

By Lemma 4 we have $v \in VP[u, y_0]$ and $x \in VP[w, y_0]$. It remains to show that $u \in VP[x, y_0]$. Suppose therefore that $w \in VP[v, y_0]$. We may suppose u, v, w, x to have been chosen to minimise $|P[v, w]|$. It follows from Claim 2 that we may now construct from $\mathcal{C}$ a new cascade $\mathcal{C}'$ by replacing C_i with the circuit

$$C'_i[x_i, v] \cup P[v, w] \cup C'_i[w, y_i] \cup A'[y_i, x_i].$$

This result contradicts the minimality of (1), and thereby completes the proof of the claim.

The same argument can be applied to AC_i -arcs which meet $A'[u_1, u_2]$. In other words, let us suppose that for some i there are two such AC_i -arcs. Let one of these arcs join vertices u and v , where $u \in VC'_i[x_i, v]$, and let the other join

vertices w and x , where $w \in VC'_i[v, y_i]$ and $x \in VC'_i[w, y_i]$. Then $x \in VA'[w, u_2]$, $u \in VA'[x, u_2]$ and $v \in VA'[u, u_2]$.

We now investigate the possible traces a cascade may have.

Claim 4: Any two consecutive digits in a trace of $\mathcal{C}$ are distinct.

Suppose that a trace of $\mathcal{C}$ contains consecutive ones. Then for some i there exist a BC_i -arc included in P and joining vertices u and v , where $u \in VC'_i[x_i, v]$, and another BC_i -arc included in P and joining vertices w and x , where $w \in VC'_i[v, y_i]$ and $x \in VC'_i[w, y_i]$, such that $C'_i[v, w]$ has no edges or internal vertices in common with the graph $G[A \cup B]$. By Claim 2 we have $x \in VP[w, y_0]$, $u \in VP[x, y_0]$ and $v \in VP[u, y_0]$. Moreover the edges of f incident on v and on w both belong to $P[v, w]$. Let

$$\begin{aligned} C &= P[v, w] \cup C'_i[w, v], \\ f^* &= f + C, \\ B^* &= B + C \\ &= A'[y_0, u_1] \cup \{e_1\} \cup A'[v_1, x_0] \cup P[x_0, w] \cup C'_i[w, v] \cup P[v, y_0] \end{aligned}$$

and

$$\begin{aligned} C_i^* &= C_i + C \\ &= C'_i[x_i, u] \cup P[u, x] \cup C'_i[x, y_i] \cup A'[y_i, x_i]. \end{aligned}$$

We now obtain a new cascade from $\mathcal{C}$ by replacing C_i with C_i^* . This result contradicts the minimality of (1).

The argument is similar if a trace of $\mathcal{C}$ contains consecutive twos. Suppose therefore that it contains consecutive zeros. Then for some i such that $1 < i < k$ the circuit C_i has a unique AC_i -arc and no BC_i -arc. The AC_i -arc joins vertices x_i and y_i . Let

$$\begin{aligned} f^* &= f + C_i, \\ A^* &= A + C_i \\ &= A'[v_1, y_i] \cup C'_i[y_i, x_i] \cup A'[x_i, u_1] \cup \{e_1\} \end{aligned}$$

and

$$C_i^* = C_{i-1} + C_i + C_{i+1}$$

$$\begin{aligned}
&= C'_{i-1}[x_{i-1}, y_{i-1}] \cup A'[y_{i-1}, x_{i+1}] \cup C'_{i+1}[x_{i+1}, y_{i+1}] \\
&\quad \cup A'[y_{i+1}, y_i] \cup C'_i[y_i, x_i] \cup A'[x_i, x_{i-1}].
\end{aligned}$$

Then f^* , A^* , B and the cascade

$$(C_1, C_2, \dots, C_{i-2}, C_i^*, C_{i+2}, C_{i+3}, \dots, C_k)$$

contradict the minimality of (1). The proof of the claim is now complete.

Claim 5: A trace of $\mathcal{C}$ contains neither 01 nor 10.

We may assume without loss of generality that a trace of $\mathcal{C}$ contains 01. Then there exists $i > 1$ such that C'_i includes a path joining x_i to a vertex $x \in VP - \{x_0, y_0\}$ and having no edges or internal vertices in common with the graph $G[A \cup B]$. By Lemma 4 the edge of f incident on x belongs to $P[x, y_0]$. Observe that the circuit

$$Y = C'_i[x_i, x] \cup P[x, y_0] \cup A'[y_0, x_i]$$

is clockwise odd under a Pfaffian orientation of K , for otherwise the choice of A and B is contradicted by A and

$$B^* = A'[v_1, x_i] \cup C'_i[x_i, x] \cup P[x, y_0] \cup A'[y_0, u_1] \cup \{e_1\}$$

since $A'[x_0, x_i] \subseteq A \cap B^*$. Similarly the circuit

$$Z = C'_i[x_i, x] \cup P[x, x_0] \cup A'[x_0, x_i]$$

is clockwise odd, as otherwise the choice of A and B is contradicted by B^* and B since $P[x, y_0] \subseteq B \cap B^*$. Thus of the three circuits Y , Z and

$$Y + Z = A + B = P \cup A'[x_0, y_0],$$

which have empty sum, only $A + B$ is clockwise even, yet an even number of them are clockwise even under a given orientation of G . This contradiction proves the claim.

By Claims 4 and 5 every second member of a trace of $\mathcal{C}$ must be 2. Having established the existence of a cascade with this property, we may now drop the requirement that A and B be chosen to maximise $|A'[v_2, y_0]| + |A'[x_0, v_1]|$. We

obtain a *reduced* trace of $\mathcal{C}$ by removing each 2 from a trace. We deal next with the case where a reduced trace has three consecutive equal digits.

Claim 6: Suppose a trace of $\mathcal{C}$ contains 02020 or 12121. Then G has an even subdivision H of $K_{3,3}$ such that $G - VH$ has a 1-factor.

Without loss of generality we may assume that a trace of $\mathcal{C}$ contains 02020. Then for some i such that $2 \leq i \leq k - 2$ the circuit C_i has a unique AC_i -arc meeting $A'[u_1, u_2]$ but it has no BC_i -arc meeting P , and a similar statement holds for C_{i+1} . The required subdivision of $K_{3,3}$ is

$$G[A'[u_1, y_0] \cup A'[x_0, v_1] \cup \{e_1\} \cup P \cup C_i \cup C_{i+1}].$$

We may henceforth assume that a reduced trace does not have three consecutive equal digits.

A set S of digits in a reduced trace is said to be *separated* if no two elements of S are consecutive. We now consider the case where a reduced trace has three separated equal digits. We shall use $*$ to denote a non-empty subsequence of a trace of $\mathcal{C}$.

Claim 7: Suppose a trace of $\mathcal{C}$ contains $12 * 212 * 21$ or $02 * 202 * 20$. Then G has an even subdivision H of $K_{3,3}$ such that $G - VH$ has a 1-factor.

Without loss of generality we may suppose that a trace of $\mathcal{C}$ contains $02 * 202 * 20$. For some h such that $2 \leq h \leq k - 2$ the path C'_h includes a subpath joining x_h to a vertex $x'_h \in VA'[u_1, u_2]$ and having no edges or internal vertices in common with the graph $G[A \cup B]$. Moreover for some i such that $h < i < k$ the path C'_{i-1} includes a subpath joining y_{i-1} to a vertex $y'_{i-1} \in VA'[u_1, x'_h]$ and having no edges or internal vertices in common with $G[A \cup B]$, and C'_i includes a subpath joining x_i to a vertex $x'_i \in VA'[u_1, y'_{i-1}]$ and having no edges or internal vertices in common with $G[A \cup B]$. Furthermore, for some j such that $i \leq j < k$ the path C'_j includes a subpath joining y_j to a vertex $y'_j \in VA'[u_1, x'_i]$ and having no edges or internal vertices in common with $G[A \cup B]$. The required subdivision of $K_{3,3}$ is

$$G[A'[u_1, y_0] \cup A'[x_h, y_j] \cup A'[x_0, v_1] \cup \{e_1\} \cup P \cup C'_h[x_h, x'_h]$$

$$\cup C'_{i-1}[y_{i-1}, y'_{i-1}] \cup C'_i[x_i, x'_i] \cup C'_j[y_j, y'_j].$$

We may henceforth assume that a reduced trace does not have three separated equal digits.

At this juncture we observe that there are only a finite number of possible traces left to investigate, because any reduced trace of length at least 9 must contain three consecutive zeros, three consecutive ones, three separated zeros or three separated ones. It can be shown that in each case there is a set of f -alternating circuits with sum $A + B$. If the clockwise parities of these circuits under a Pfaffian orientation of K do not yield a contradiction, then the fact that $A + B$ is clockwise even under a Pfaffian orientation of K implies, by Lemma 2, that exactly one of A and B is clockwise even. In this case G must be non-Pfaffian, as it cannot be oriented so that all the f -alternating circuits are clockwise odd. In such graphs G we therefore search for a subgraph H , reducible to an even subdivision of $K_{3,3}$, such that $G - VH$ has a 1-factor.

The next claim is proved by investigating the condition under which a contradiction can be derived from the clockwise parities, under a Pfaffian orientation of K , of a set of f -alternating circuits with sum $A + B$.

Claim 8: A reduced trace of $\mathcal{C}$ is of odd length.

Suppose that a reduced trace of $\mathcal{C}$ is of even length. First we construct a cascade $\mathcal{D}$ along P from y_0 to x_0 . If a reduced trace has no 1, then $\mathcal{D}$ consists only of the circuit

$$D_1 = X \cup A'[y_k, x_0] \cup P \cup A'[y_0, x_1].$$

Otherwise there exists a vertex $z_1 \in VX \cap VP$. We may assume z_1 chosen to minimise $|X[w_1, z_1]|$, where $w_1 = x_1$. In this case we define

$$D_1 = X[w_1, z_1] \cup P[z_1, y_0] \cup A'[y_0, w_1].$$

We also let w_2 be the vertex of $VX \cap (VP[z_1, z_0] - \{z_0\})$ that minimises $|P[w_2, z_0]|$, where $z_0 = y_0$.

If there exists a vertex $z_2 \in (VX[w_2, y_k] - \{w_2\}) \cap VP$, then we repeat the argument. Thus we may assume z_2 chosen to minimise $|X[w_2, z_2]|$. We then define

$$D_2 = X[w_2, z_2] \cup P[z_2, w_2].$$

We also let w_3 be the vertex of $VX \cap (VP[z_2, z_1] - \{z_1\})$ that minimises $|P[w_3, z_1]|$.

If there exists a vertex $z_3 \in (VX[w_3, y_k] - \{w_3\}) \cap VP$, then we repeat the argument inductively. Thus there exists $l > 0$ such that

$$(VX[w_l, y_k] - \{w_l\}) \cap VP = \emptyset.$$

We define

$$D_l = X[w_l, y_k] \cup A'[y_k, x_0] \cup P[x_0, w_l].$$

The required cascade $\mathcal{D}$ is then $(D_1, D_2, \dots, D_l)$.

We observe that the number of ones in a trace of $\mathcal{C}$ is $l - 1$. Similarly the number of zeros is $k - 1$. Hence $k + l$ is even by assumption. Moreover

$$\sum_{i=1}^k C_i + \sum_{j=1}^l D_j = A + B.$$

(For any i each member of a PD_i -arc belongs to one circuit in $\mathcal{C}$ and two circuits in $\mathcal{D}$. The remaining elements of P belong to just one circuit in $\mathcal{D}$ and to no circuits in $\mathcal{C}$. Similar results hold for the edges of $A'[x_0, y_0]$. The remaining edges of $X \cup A'[y_0, x_1] \cup A'[y_k, x_0]$ belong to one circuit in each of $\mathcal{C}$ and $\mathcal{D}$.) In addition, it is easy to see that K may be oriented so that the circuits in $\mathcal{C}$ and $\mathcal{D}$ are directed. For example, if $\{M, N\}$ is a bipartition of K , we may orient the edges of f from M to N and the remaining edges from N to M . Under this orientation the circuit $A + B$ is clockwise odd. As it is the only clockwise odd circuit in the set

$$S = \{C_1, C_2, \dots, C_k, D_1, D_2, \dots, D_l, A + B\},$$

and S has empty sum and odd cardinality, we now have a contradiction because under a Pfaffian orientation of K the circuit $A + B$ is the only clockwise even member of S . This contradiction proves the claim.

In all the remaining cases we must find a subgraph H of G , reducible to an even subdivision of $K_{3,3}$, such that $G - VH$ has a 1-factor. We simplify the work by establishing two further claims which enable us to manipulate a trace of $\mathcal{C}$.

Claim 9: Let G' be a graph with a subgraph H' , reducible to an even subdivision of $K_{3,3}$, such that $G' - VH'$ has a 1-factor. Suppose that G' has a cascade

$\mathcal{C}'$ with a trace whose first element is 0 or 1. Let G have a cascade $\mathcal{C}$ with a trace consisting of 2 followed by the trace of $\mathcal{C}'$. Then G has a subgraph H , reducible to an even subdivision of $K_{3,3}$, such that $G - VH$ has a 1-factor.

The path C'_1 includes a subpath joining x_1 to a vertex $x'_1 \in VA'[u_1, u_2]$ and having no edges or internal vertices in common with $G[A \cup B]$. Contraction of the circuit $C'_1[x_1, x'_1] \cup A'[x'_1, x_1]$ yields the graph G' . Since G' has a subgraph H' , reducible to an even subdivision of $K_{3,3}$, such that $G' - VH'$ has a 1-factor, the claim follows.

A corresponding result holds if the last element of a trace of $\mathcal{C}'$ is 0 or 1 and a trace of $\mathcal{C}$ consists of that of $\mathcal{C}'$ followed by 2. We may therefore assume that neither the first nor the last element of a trace of $\mathcal{C}$ is 2.

Claim 10: Suppose that the first element of a trace of $\mathcal{C}$ is 0. Then $\mathcal{C}$ has another trace obtained by replacing the first 0 with 1.

Note first that the path $C'_1[x_1, y_1]$ has no edges or internal vertices in common with $G[A \cup B]$. Let

$$\begin{aligned} f^* &= f + C_1, \\ A^* &= A + C_1 \\ &= A'[u_1, x_1] \cup C'_1[x_1, y_1] \cup A'[y_1, v_1] \cup \{e_1\} \end{aligned}$$

and

$$\begin{aligned} B^* &= B + C_1 \\ &= P \cup A'[y_0, y_1] \cup C'_1[y_1, x_1] \cup A'[x_1, u_1] \cup \{e_1\} \cup A'[v_1, x_0]. \end{aligned}$$

The required trace is calculated by replacing A and B with the f^* -alternating circuits A^* and B^* respectively.

A corresponding result holds if the first element of a trace of $\mathcal{C}$ is 1. We also obtain corresponding results for the last element of a trace of $\mathcal{C}$.

Claim 11: A reduced trace of $\mathcal{C}$ has length 1.

Suppose first that a reduced trace of $\mathcal{C}$ has length at least 5. By Claim 10 we may assume that the first and last digits are 0. By Claim 7 the third digit must therefore be 1. However, by Claim 10 there is another reduced trace of $\mathcal{C}$

obtained by replacing the first and last digits with 1, in violation of Claim 7. The argument is similar if a reduced trace of $\mathcal{C}$ has length 3, except that Claim 6 is used instead of Claim 7. Claim 11 now follows from Claim 8.

Thus $\mathcal{C}$ has a reduced trace of length 1. By symmetry and Claim 9, we may assume that 0 is a trace of $\mathcal{C}$. Then

$$G[A \cup B \cup C_1 \cup C_2]$$

is the required subdivision of $K_{3,3}$. $\square$

Theorem 5 *Let G be a 1-extendible graph that can be obtained from a 1-extendible bipartite graph K by a 2-ear adjunction. Suppose that $G[C \cup D]$ is bipartite whenever C and D are alternating circuits that traverse both ears and have the property that there are just two $\bar{C}D$ -arcs. Then G is non-Pfaffian if and only if it has a subgraph H , reducible to an even subdivision of $K_{3,3}$, such that $G - VH$ has a 1-factor.*

Proof. We have already seen that if G has H as a subgraph then G is non-Pfaffian. Suppose therefore that G is non-Pfaffian. Suppose also that G is formed from a 1-extendible bipartite graph K by the adjunction of ears E_1 and E_2 . If K is non-Pfaffian, then K has the required subgraph H by Theorem 2, and therefore so does G . We may suppose therefore that K is Pfaffian.

Suppose that $G - E_1$ is 1-extendible. Since $G - E_1$ is obtained from the bipartite graph K by the adjunction of the single ear E_2 , it follows that $G - E_1$ is also bipartite. Since G is obtained from $G - E_1$ by the adjunction of a single ear, G is also bipartite. Therefore G has the required subgraph by Theorem 2. The argument is similar if $G - E_2$ is 1-extendible.

We may now assume that neither $G - E_1$ nor $G - E_2$ is 1-extendible. Let E_1 join vertices u_1 and v_1 and fix a 1-factor f of K . Since K is 1-extendible, it has a path P , joining u_1 and v_1 , such that each vertex of $VP - \{u_1\}$ is incident with an edge of $P \cap f$. Thus $P \cup E_1$ is an alternating circuit in $G - E_2$ if $G - E_2$ is bipartite, and in this case we reach the contradiction that $G - E_2$ is 1-extendible. Hence $G - E_2$ is not bipartite. Similarly $G - E_1$ is not bipartite.

Next, we may clearly assume without loss of generality that E_1 and E_2 both have cardinality 1. Therefore we may let $E_1 = \{e_1\}$ and $E_2 = \{e_2\}$. Thus e_1 and e_2 are distinct edges of $EG - f$ such that neither $G - \{e_1\}$ nor $G - \{e_2\}$ is bipartite and neither is 1-extendible, but $G - \{e_1, e_2\}$ is bipartite and 1-extendible. As G is 1-extendible, there must be an f -alternating circuit X containing e_1 . Since $G - \{e_2\}$ is not 1-extendible, X must also contain e_2 . It follows that e_1 and e_2 are independent edges, as X is f -alternating.

Since K is Pfaffian we may assume given a Pfaffian orientation of K . Extend this orientation to G by orienting e_1 and e_2 arbitrarily. As G is not Pfaffian, there must exist a clockwise even alternating circuit A in G . Recalling that some 1-factor in G must have its sign opposite to that of f , we find that A may be chosen to be f -alternating. Being clockwise even, it cannot be alternating in K , and therefore cannot be a circuit of K . Thus A contains e_1 or e_2 and hence both. There must also be a clockwise odd alternating circuit B containing e_1 and e_2 for otherwise a Pfaffian orientation of G could be realised by reorienting e_1 or e_2 . We can also choose B to be f -alternating. By Lemma 3 we find that f , A and B may be chosen so that there are just one or two AB -arcs, each containing e_1 or e_2 and their union containing both.

If there are two AB -arcs, then $G[A \cup B]$ is bipartite by hypothesis. However one of these arcs must contain e_1 and the other e_2 . Let $A' = A - \{e_1\}$, and let e_2 join vertices u_2 and v_2 , where $e_2 \in A'[u_1, v_2]$. If there exists an $\bar{A}B$ -arc Q joining a vertex $x \in VA'[u_1, u_2]$ to a vertex $y \in VA'[v_1, v_2]$, then since $G[A \cup B]$ is bipartite we find that either

$$Q \cup A'[y, x]$$

or

$$Q \cup A'[x, u_1] \cup \{e_1\} \cup A'[v_1, y]$$

is an f -alternating circuit containing just one member of $\{e_1, e_2\}$. From this contradiction to the fact that neither $G - \{e_1\}$ nor $G - \{e_2\}$ is 1-extendible, together with the knowledge that one AB -arc contains e_1 and the other e_2 , we deduce that $A + B$ is the union of two disjoint circuits.

By Lemma 2 one of these circuits is a clockwise even alternating circuit. It is therefore possible to modify B so that there is only one AB -arc. Then $A + B$ is

a clockwise even alternating circuit. We can now invoke Theorem 4 to draw the desired conclusion. $\square$

In order to complete the analysis of a 2-ear adjunction to a bipartite graph, it remains to consider the case where there are two AB -arcs. This case is considerably more complicated, and is currently being studied by the authors.

References

- [1] M. H. Carvalho, C.L. Lucchesi and U.S.R. Murty. Ear decompositions of matching covered graphs. Technical report, University of Waterloo, 1998.
- [2] J. Edmonds, L. Lovász and W.R. Pulleyblank. Brick decompositions and the matching rank of graphs. *Combinatorica*, 2:247-274, 1982.
- [3] P.W. Kasteleyn. Graph theory and crystal physics. In F. Harary, editor, *Graph theory and theoretical physics*, pages 43-110. Acad. Press, New York, 1967.
- [4] C.H.C. Little. Kasteleyn's theorem and arbitrary graphs. *Canad. J. Math.*, 25:758-764, 1973.
- [5] C.H.C. Little. A characterization of convertible $(0,1)$ -matrices. *Journal of Combinatorial Theory (B)*, 18:187-208, 1975.
- [6] C.H.C. Little and F. Rendl. Operations preserving the Pfaffian property of a graph. *J. Australian Math. Soc. (Series A)*, 50:248-257, 1991.
- [7] L. Lovász. Matching structure and the matching lattice. *Journal of Combinatorial Theory (B)*, 43:187-222, 1987.
- [8] L. Lovász and M.D. Plummer. *Matching Theory*, Akadémiai Kiadó, Budapest, 1986.